

Culver School District Protocol and Procedures for Management of Sports-Related Concussions

Medical management of a sports-related concussion is evolving. In recent years, there has been a significant amount of research into sports-related concussions in middle school/high school athletes and legislation has been adopted to protect student-athletes and to ensure best practices among Oregon school districts.

Culver School District (also referred to as the "District") has established this Protocol and Procedures for Management of Sports-Related Concussions (this "Protocol") to educate and guide persons who instruct or train members of a school athletic team, including employees, volunteers, and contract coaches/instructors ("Coaches"), in the treatment and management of sports-related concussions and to comply with applicable laws. This Protocol outlines procedures for staff to follow in managing head injuries and outlines school policy as it pertains to a student-athlete's resumption of athletic activities, including practice or conditioning, following a concussion ("Return to Play").

Culver School District seeks to provide a safe "Return to Play" for all athletes after injury, particularly after a concussion. In order to effectively and consistently manage concussions, procedures have been developed to aid in insuring that concussed athletes are identified, treated, referred appropriately, receive proper follow-up medical care during the school day, including academic assistance, and are fully recovered prior to a "Return to Play".

In addition to recent research, ORS 336.485, OAR 581-022-0421, and five (5) primary documents were consulted in developing this protocol.

- 1) Journal of Athletic Training 2013;48(4):554–575 doi: 10.4085/1062-6050-48.4.05 by the National Athletic Trainers' Association, Inc www.natajournals.org consensus statement Consensus Statement on Concussion in Sport: The 4th International Conference on Concussion in Sport, Zurich, November 2012.
- 2) Journal of Athletic Training 2014;49(2):245–265 doi: 10.4085/1062-6050-49.1.07 by the National Athletic Trainers' Association, Inc www.natajournals.org position statement National Athletic Trainers' Association Position Statement: Management of Sport Concussion and
- 3) Max's Law: Concussion Management Implementation Guide for School Administrators ("Max's Law").
- 4) CBIRT/OCAMP sample school district concussion protocol
- 5) Silverberg ND, Iverson GL. Is Rest After Concussion "The Best Medicine?": Recommendations for Activity Resumption Following Concussion in Athletes, Civilians, and Military Service Members. J Head Trauma Rehabil. 2013; 28 (4):250-259.

The Team members who provided input that led to the development of this Protocol are; {000131470-00481657;3}

- CULVER Athletic Trainer- Courtney Miller
- CULVER HS Athletic Director- Shea Little
- CULVER Nurse- Kara Nielsen
- Medical Director of The Center Foundation Dr. Viviane Ugalde
- ATC Concussion Coordinator, The Center Foundation Lindsay Hagler
- ATC Rural Outreach Coordinator, The Center Foundation- Emily Schleicher
- High Desert ESD Concussion Lesion Sue Hayes

Contents:

- I. Recognition of Concussion
- II. Management and Referral Guidelines for All Staff
- III. Procedures for the Certified Athletic Trainer (AT)
- IV. Guidelines and Procedures for Coaches
- V. Follow-up care during the School Day
- VI. Return to Play Procedure
- VII. Protocol Updates

I. Recognition of Concussion

- A. For purposes of this Protocol, common signs and symptoms of sports-related concussions ("Signs and Symptoms" or "Symptomatic") include:
 - 1. Signs (observed by others):
 - Athlete appears dazed
 - Confusion (about assignment, plays, days of week etc.)
 - Forgets plays or instructions
 - Unsure about game, score, opponent
 - Moves clumsily (altered coordination)
 - Balance problems
 - Personality change
 - Responds slowly to questions
 - Forgets events prior to hit
 - Forgets events after the hit
 - Loss of consciousness ("LOC") for any duration
 - 2. Symptoms (reported by athlete):
 - Headache
 - Fatigue
 - Nausea or vomiting
 - Double vision, blurry vision
 - Sensitive to light or noise/ringing in ears
 - Feels sluggish
 - Feels "foggy"
 - Problems concentrating
 - Problems remembering

- 3. These signs and symptoms identified above are indicative of probable concussion. Other causes or symptoms should also be considered.
- B. Cognitive Impairment Testing (altered or diminished cognitive function)

General cognitive status can be determined by simple sideline cognitive testing with SCAT5 (Form A SCAT5) If an AT is present then the AT will do a sideline assessment. If no AT is present then the coach can perform the testing using the Coaches Report (Form B-Coaches/CMT Concussion Report Form).

- C. Neurocognitive testing requirements
 - 1. Neurocognitive testing (ImPACT) is a research-based software tool utilized to evaluate recovery after concussion. This testing evaluates multiple aspects of neurocognitive function, including memory, attention, and brain processing speed, reaction time, and post-concussion symptoms.
 - a. Neurocognitive testing shall be utilized to help determine recovery after concussion. This test is given by the AT or a practitioner that is trained in ImPACT.
- D. Athletes participating in athletics will be offered a baseline test prior to participating in sports during grades 9-12. These tests are administered by The Center Foundation Athletic Trainer.
- E. Due to growth and brain development, baselines are obtained every 2 years, typically in the freshman and junior years. All athletes in their junior year will be required to take a "new" baseline test prior to participating in Athletics.
- F. Post-concussion neurocognitive testing is performed to help determine concussion recovery and is done at the school by athletic trainer at no charge or at The Center as part of concussion recovery assessment. Comparisons of results are made to baseline testing or if baselines aren't available, then to age matched controls

II. Management and Referral Guidelines for All Staff

- A. Guidelines for Responding to this Protocol
 - 1. Athletes experiencing/exhibiting signs and symptoms of concussion will be removed from participation and shall be evaluated by the athletic trainer or concussion team member and then referred to primary care providers, or emergency room (Form A-SCAT5 or B-Coaches/CMT Concussion Report Form).
 - 2. Athletes experiencing a witnessed loss of consciousness of any duration should be transported immediately to the nearest hospital emergency department via emergency vehicle.
 - 3. Any athlete who has signs and symptoms, and who is not stable (i.e., condition is persisting or deteriorating), must be transported immediately to the nearest hospital emergency department via emergency vehicle.

- 4. An athlete who is symptomatic, but stable, may be transported by his/her parents. The parents should be advised to contact a licensed physician, physician's assistant, Nurse practitioner, psychologist, Desert Orthopedic Center, The Center Now Care Clinic or seek care at the nearest hospital emergency department (collectively, "Health Care Professional"). The parents will be provided an informational sheet regarding concussion (Form C-Parent Concussion Information Packet).
 - a. ALWAYS advise parents the option of emergency transportation, even if you do not feel it is necessary.

III. Procedures for the Certified Athletic Trainer – This section applies to schools with AT recourses

- A. Following a suspected concussion, the Oregon Health Licensing Board requires the Certified Athletic Trainer at the athlete's school (the "AT") to assess the injury, or provide guidance to the Coach(s) of the sport the athlete is currently participating in ("Sport Coach") if unable to personally attend to the athlete.
 - 1. The AT will perform serial assessments following recommendations in the NATA Position Statement SCAT5 assessment tool (Form A-SCAT5)
 - a. The AT will notify the athlete's parents and give written and verbal home and follow-up care instructions.
 - 2. Referral to a health care professional will be mandatory and made when medically appropriate.
- B. The AT will notify the Nurse or concussion management team (CMT) Team at the athlete's school of the injury, as soon as possible, so they can initiate appropriate follow-up care with the concussion management team upon the athlete's return to school.
 - 1. The AT will continue to provide coordinated care with the School Nurse or CMT Team for the duration of the injury. Care of athletes with persistent symptoms will be managed in conjunction with the athlete's physician.
 - 2. The School Nurse or CMT Team will communicate with the athlete's guidance counselor regarding the athlete's neurocognitive and recovery status, if needed.
- C. The AT is responsible for administering ImPACT testing.
 - 1. The post-concussion ImPACT test will be performed within 72 hours if there is a question regarding diagnosis of concussion. Otherwise, the post injury ImPACT test will be performed at the time of athlete's symptoms resolve.
 - 2. The AT will review the post-concussion test data with the athlete and the athlete's parent.
 - 3. The AT will forward testing results to the athlete's Health Care Professional, with parental permission and a signed release of medical information form.

- 4. The AT will advise and regularly update the Sport Coach, including the Sport Coach in the following season if the athlete transitions between sport seasons during the course of the Progression (defined and described in Section VI below), regarding the athlete's participation limitations.
- 5. The AT will monitor the athlete, and keep the School Nurse or CMT Team informed of the individual's symptomatology and neurocognitive status, for the purposes of developing or modifying the Health Care Plan, as defined below, for the student-athlete. Athletes and parents sign a release for treatment and coordination of care to include school Nurse, CMT Team, and administrators as a part of their sport packet before playing sports.
- 6. The AT will work with the Health Care Professional to coordinate the progression.
- 7. The AT will maintain appropriate computerized documentation regarding assessment and management of the injury.

IV. Procedures for Coaches:

- A. All Coaches shall receive annual training (no less than once every twelve months), prior to initiation of the season for the sport in which that Coach instructs or trains, to learn how to recognize the symptoms of a concussion. Each school in the District that sponsors athletics shall annually develop a list of all Coaches, identify the resources to be used to provide the training, develop training timelines for all Coaches, and document that each Coach completes the training described in subsection (B) below. Training will be tracked and documented annually.
- B. Annual training shall include training on the following topics:
 - (a) Training in how to recognize the signs and symptoms of a concussion;
 - (b) Training in strategies to reduce the risk of concussions;
 - (c) Training in how to seek proper medical treatment for a person suspected of having a concussion; and
 - (d) Training in procedures of how an athlete may safely return to participation.

When a concussion is suspected, Coaches shall follow the general principles of RECOGNIZE, REMOVE, and REFER

- C. Recognize concussion signs and symptoms (Form B-Coaches/CMT Concussion Report Form)
 - 1. Use of Coaches Report Form (Form B-Coaches/CMT Concussion Report Form) to record signs and symptoms, copy should be given to parent/guardian at time of the incident.
- D. Remove from activity

- 1. If a Coach suspects the athlete has sustained a concussion, the athlete shall be removed from activity immediately and for the day.
 - a. Any athlete who exhibits signs and symptoms following an observed or suspected blow to the head or body will be removed immediately from participation, assessed, and will not be allowed to Return to Participation that day.

E. Refer the athlete for medical evaluation

- 1. Coaches shall report all head injuries to the AT or CMT Team, as soon as possible, for medical assessment and management, for coordination of home instructions and for follow-up care.
- 2. Coaches should seek assistance from the host site AT if at an away contest.
- 3. If the AT is unavailable, or the athlete is injured at an away event, The Sport Coach is responsible for:
 - a. Contacting the athlete's parents to inform them of the injury and to make arrangements for them to pick-up the athlete.
 - b. Providing the AT and CMT Team with the athlete's name and home phone number, so that the AT or CMT Team can initiate follow-up. Additional copies are available from the AT.
 - c. Reminding the athlete to report directly to the School Nurse or CMT Team before school starts on the day the student returns to school after the injury.
- 4. In the event that an athlete's parents cannot be reached, and the athlete is able to be sent home:
 - a. The AT or Sport Coach should ensure that the athlete will be with a responsible individual, who is capable of monitoring the athlete and understanding the home care instructions, before allowing the athlete to go home.
 - b. The AT or Sport Coach should continue efforts to reach the parent.
 - c. If there is any question about the status of the athlete, or if the athlete is not able to be monitored appropriately, the athlete should be referred to the emergency department at the nearest hospital for evaluation. The Sport Coach or AT should accompany the athlete and remain with the athlete until the athlete's parents arrive.
 - d. Athletes exhibiting Signs and Symptoms should not be permitted to drive home.

V. FOLLOW-UP CARE OF THE ATHLETE DURING THE SCHOOL DAY

- A. Responsibilities of the School Nurse or CMT Team after notification of student's suspected concussion.
 - 1. The athlete will be instructed to report to the School Nurse or CMT Team for Return to Learn procedures.
 - 2. Immediately notify the student's guidance counselor and teachers of the injury that a Concussion Accommodation Plan has been developed.
 - 3. Notify the student's physical education teacher immediately that the athlete is restricted from physical activity until further notice from the School Nurse or CMT Team.
 - 4. Bed rest should be no more than 3 days. Athletes should return to light activity following concussion guidelines. The school Nurse or CMT Team will utilize (Form D-OSAA Concussion Return Form) to assist with progression and direction from the physician.
 - 5. If the School Nurse or CMT Team receives notification of a student-athlete who has sustained a concussion from someone other than the AT (i.e. the athlete, athlete's parent, Sport Coach, physician, etc.), the AT should be notified when AT resources are available.
 - 6. Monitor the athlete as needed during recovery.
- B. Responsibilities of the student's guidance counselor
 - 1. Monitor the student closely and recommend appropriate academic accommodations for students who are exhibiting post-concussion symptoms. Reference materials for concussion related academic accommodations are listed in (Form D-OSAA Concussion Return Form).
 - 2. Communicate with the School Nurse or CMT Team on a regular basis, to provide the most effective care for the student.

VI. RETURN TO PLAY PROCEDURES AFTER CONCUSSION

- A. Returning to Activity on the same day of injury
 - 1. An athlete who exhibits Signs and Symptoms following an observed or suspected blow to the head or body, or is otherwise diagnosed with a concussion is not permitted to Return to Play on the day of the injury.
 - 2. "When in doubt, hold them out."
- B. Return to Play after suspected concussion
 - 1. Following a concussion, athletes will not be permitted to Return to Play until the athlete has completed the stepwise progression outlined in this subsection (Form D-Concussion Return Form).

- 2. As described in the Zurich 2012 the progression consists of the following steps:
 - a. No activity on day of concussion.
 - b. No bed rest after 1-3 days max.
 - c. Allowed light aerobic exercise walking, stationary bike 0 3 days
 - d. Sport-specific training (e.g., skating in hockey, running in soccer)
 - e. Non-contact training drills
 - f. Full-contact training after medical clearance
 - g. Game play (unrestricted activity)

Return to play progression will be monitored by the AT. In schools without AT's, then parents, or school Nurse or CMT Team members will be provided with (Form D-OSAA Concussion Return Form) for progression.

- 3. The athlete must meet all of the following criteria in each step of the Progression in order to return to participation:
 - a. No longer exhibits signs, symptoms or behaviors consistent with a concussion at rest and with exertion (including mental exertion in school);
 - b. Is participating in full school hours and classroom activities without accommodations, except for the need for more time for makeup work.
 - c. If athlete has a valid baseline test, is within normal range of baseline on post-concussion neurocognitive testing
 - d. If athlete has no baseline, then is testing within a range consistent with their academic performance and compared to age matched controls.
 - e. Have written clearance from the athlete's Health Care Professional using the OSAA form (Form D-OSAA Concussion Return Form)
- 4. The school Nurse or CMT Team will supervise the Return to Learn and determine the athlete's status in the progression with physician recommendations.
- 5. The AT or CMT Team will supervise the Return to Play and determine the athlete's status in the progression with physician recommendations.
- 6. The AT or CMT Team and athlete will discuss appropriate activities for each day the athlete participates in high school athletics. The athlete's participation will be limited to those appropriate activities until the AT or CMT Team instructs otherwise.
- 7. The athlete should see the school Nurse or CMT Team or counselor as needed for academic reassessment and instructions until he/she has progressed to the Return to Play progression.
- 8. The athlete should see the AT or CMT Team member as needed for re-assessment and instructions until he/she has progressed to unrestricted activity, and received written clearance for Return to Play.
- 9. No additional testing is needed once the athlete is cleared to play.

VII. PROTOCOL UPDATES

(000404470 00404577 0)		
{000131470-00481657;3}		

A. Given that concussion related knowledge and best practices are rapidly evolving, Culver School District will periodically re-evaluate and update the Protocol. It is recommended we review the protocol every 1 year on May 1, for completion by June 1.

Appendix A SCAT5

Downloaded from http://bjsm.bmj.com/ on May 12, 2017 - Published by group.bmj.com
BJSM Online First, published on April 26, 2017 as 10.1136/bjsports-2017-097506SCAT5

To download a clean version of the SCAT tools please visit the journal online (http://dx.dol.org/10.1136/bjsports-2017-097506SCATS)



SPORT CONCUSSION ASSESSMENT TOOL - 5TH EDITION

FOR USE BY MEDICAL PROFESSIONALS ONLY

supported by









Patient details		
Name:		
Examiner:		
Date of Injury:	Time:	

WHAT IS THE SCAT5?

The SCAT5 is a standardized tool for evaluating concussions designed for use by physicians and licensed healthcare professionals1. The SCAT5 cannot be performed correctly in less than 10 minutes.

If you are not a physician or licensed healthcare professional, please use the Concussion Recognition Tool 5 (CRT5). The SCAT5 is to be used for evaluating athletes aged 13 years and older. For children aged 12 years or younger, please use the Child SCAT5.

Preseason SCAT5 baseline testing can be useful for interpreting post-injury test scores, but is not required for that purpose. Detailed instructions for use of the SCAT5 are provided on page 7. Please read through these instructions carefully before testing the athlete. Brief verbal instructions for each test are given in italics. The only equipment required for the tester is a watch or timer.

This tool may be freely copied in its current form for distribution to individuals, teams, groups and organizations. It should not be altered in any way, re-branded or sold for commercial gain. Any revision, translation or reproduction in a digital form requires specific approval by the Concussion in Sport Group.

Recognise and Remove

A head impact by either a direct blow or indirect transmission of force can be associated with a serious and potentially fatal brain injury. If there are significant concerns, including any of the red flags listed in Box 1, then activation of emergency procedures and urgent transport to the nearest hospital should be arranged.

Key points

- Any athlete with suspected concussion should be REMOVED FROM PLAY, medically assessed and monitored for deterioration. No athlete diagnosed with concussion should be returned to play on the day of injury.
- · If an athlete is suspected of having a concussion and medical personnel are not immediately available, the athlete should be referred to a medical facility for urgent assessment
- · Athletes with suspected concussion should not drink alcohol, use recreational drugs and should not drive a motor vehicle until cleared to do so by a medical professional.
- · Concussion signs and symptoms evolve over time and it is important to consider repeat evaluation in the assessment of concussion.
- · The diagnosis of a concussion is a clinical judgment, made by a medical professional. The SCAT5 should NOT be used by itself to make, or exclude, the diagnosis of concussion. An athlete may have a concussion even if their SCAT5 is "normal".

Remember:

- The basic principles of first aid (danger, response, airway, breathing, circulation) should be followed.
- Do not attempt to move the athlete (other than that required for airway management) unless trained to do so.
- · Assessment for a spinal cord injury is a critical part of the initial on-field assessment.
- Do not remove a helmet or any other equipment unless trained to do so safely.

Concussion in Sport Group 2017

Davis GA, et al. Br J Sports Med 2017;0:1-8. doi:10.1136/bjsports-2017-0975065CAT5

Copyright Article author (or their employer) 2017. Produced by BMJ Publishing Group Ltd under licence.

IMMEDIATE OR ON-FIELD ASSESSMENT

The following elements should be assessed for all athletes who are suspected of having a concussion prior to proceeding to the neurocognitive assessment and ideally should be done on-field after the first first aid / emergency care priorities are completed.

If any of the "Red Flags" or observable signs are noted after a direct or indirect blow to the head, the athlete should be immediately and safely removed from participation and evaluated by a physician or licensed healthcare professional.

Consideration of transportation to a medical facility should be at the discretion of the physician or licensed healthcare professional.

The GCS is important as a standard measure for all patients and can be done serially if necessary in the event of deterioration in conscious state. The Maddocks questions and cervical spine exam are critical steps of the immediate assessment; however, these do not need to be done serially.

STEP 1: RED FLAGS

RED FLAGS:

- Neck pain or tenderness
- Double vision
- Weakness or tingling/ burning in arms or legs . Vomiting
- Severe or increasing
- · Seizure or convulsion
 - Loss of consciousness
 - Deteriorating conscious state

 - Increasingly restless, agitated or combative

STEP 2: OBSERVABLE SIGNS

Witnessed Observed on Video		
Lying motionless on the playing surface	Y	N
Balance / gait difficulties / motor incoordination: stumbling, slow / laboured movements	Y	N
Discrientation or confusion, or an inability to respond appropriately to questions	Y	N
Blank or vacant look	Y	N
Facial injury after head traums	Y	N

STEP 3: MEMORY ASSESSMENT MADDOCKS QUESTIONS²

"I am going to sak you a few questions, please listen carefully and give your best effort. First, tell me what happened?"

Mark Y for correct answer / N for Incorrect		
What venue are we at today?	Y	N
Which half is it now?	Y	N
Who scored last in this match?	Y	N
What team did you play last week / game?	Y	N
Did your team win the last game?	Y	N

Note: Appropriate sport-specific questions may be substituted.

Name:		_
DOB:		_
Address:		_
ID number:		_
Examiner:		_
		_

STEP 4: EXAMINATION GLASGOW COMA SCALE (GCS)³

Time of assessment			
Date of assessment			
Best eye response (E)			
No eye opening	1	1	1
Eye opening in response to pain	2	2	2
Eye opening to speech	3	3	3
Eyes opening apontaneously	4	4	4
Best verbal response (V)			
No verbal response	1	1	1
Incomprehensible sounds	2	2	2
Inappropriate words	3	3	3
Confused	4	4	4
Oriented	5	5	5
Best motor response (M)			
No motor response	1	1	1
Extension to pain	2	2	2
Abnormal flexion to pain	3	3	3
Flexion / Withdrawal to pain	4	4	4
Localizes to pain	5	5	5
Obeys commands	6	6	6
Glasgow Coma score (E + V + M)			

CERVICAL SPINE ASSESSMENT

Does the athlete report that their neck is pain free at rest?	Y	N
If there is NO neck pain at rest, does the athlete have a full range of ACTIVE pain free movement?	Y	N
is the limb strength and sensation normal?	Y	N

In a patient who is not lucid or fully conscious, a cervical spine injury should be assumed until proven otherwise.

Concussion in Sport Group 2017

OFFICE OR OFF-FIELD ASSESSMENT

Please note that the neurocognitive assessment should be done in a distraction-free environment with the athlete in a resting state.

STEP 1: ATHLETE BACKGROUND

Sport / team / school:_

Date / time of injury:		
Years of education completed:		
Age:		
Gender: M / F / Other		
Dominant hand: left / neither / right		
How many diagnosed concussions has the athlete had in the past?:		
When was the most recent concussion?:		
How long was the recovery (time to being cleared to pli from the most recent concussion?:	ay)	_(days)
Has the athlete ever been:		_,,,,
Hospitalized for a head injury?	Yes	No
Diagnosed / treated for headache disorder or migraines?	Yes	No
Diagnosed with a learning disability / dyslexia?	Yes	No
Diagnosed with ADD / ADHD?	Yes	No
Diagnosed with depression, anxiety or other psychiatric disorder?	Yes	No
Current medications? If yes, please list:		

Name: DOB:		
Address:		
ID number:		
Examiner:		
Date:		

2

STEP 2: SYMPTOM EVALUATION

The athlete should be given the symptom form and asked to read this instruction paragraph out loud then complete the symptom scale. For the baseline excessment, the athlete should rate thin/her symptoms based on how he/she typically feels and for the post injury assessment the athlete should rate their symptoms at this point in time.

Please Check:

Baseline

Post-Injury

Please hand the form to the athlete

	none	-	lid	mod	erate	3 61	979
Headache	0	1	2	3	4	5	6
"Pressure in head"	0	1	2	3	4	5	6
Neck Pain	0	1	2	3	4	5	6
Nauses or vomiting	0	1	2	3	4	5	6
Dizziness	0	1	2	3	4	5	6
Blumed vision	0	1	2	3	4	5	6
Balance problems	0	1	2	3	4	5	6
Sensitivity to light	0	1	2	3	4	5	6
Sensitivity to noise	0	1	2	3	4	5	6
Feeling slowed down	0	1	2	3	4	5	6
Feeling like "in a fog"	0	1	2	3	4	5	6
"Don't feel right"	0	1	2	3	4	5	6
Difficulty concentrating	0	1	2	3	4	5	6
Difficulty remembering	0	1	2	3	4	5	6
Fatigue or low energy	0	1	2	3	4	5	6
Confusion	0	1	2	3	4	5	6
Drowsiness	0	1	2	3	4	5	6
More emotional	0	1	2	3	4	5	6
Irritability	0	1	2	3	4	5	6
Sadness	0	1	2	3	4	5	6
Nervous or Anxious	0	1	2	3	4	5	6
Trouble falling saleep (if applicable)	0	1	2	3	4	5	6
Total number of symptoms:							f 22
Symptom severity score:						af	132
Do your symptoms get worse with	physic	al activ	rity?		1	Y N	
Do your symptoms get worse with	menta	ectivi	ty?		1	Y N	
If 100% is feeling perfectly norms percent of normal do you feel?	l, what						
If not 100%, why?							

Please hand form back to examiner

of 5

of 15

Score (of 10)

STEP 3: COGNITIVE SCREENING Standardised Assessment of Concussion (SAC)4 ORIENTATION What month is it? What is the date today? What is the day of the week? What year is it? What time is it right now? (within 1 hour) Orientation acore IMMEDIATE MEMORY The Immediate Memory component can be completed using the traditional 5-word per trial list or optionally using 10-words per trial to minimise any ceiling effect. All 3 trials must be administered irrespective of the number correct on the first trial. Administer at the rate of one word per second. Please choose EITHER the 5 or 10 word list groups and circle the specific word list chosen for this test. I am going to test your memory. I will read you a list of words and when I am done, repeat back as many words as you can remember, in any order. For Trials 2.6.3: I am going to repeat the same list again. Repeat back as many words as you can remember in any order, even if you said the word before. Scare (of 5) List Alternate 5 word lists Trial 1 Trial 2 Trial 3 A Finger Blanket Insect Penny Lemon В Candle Paper Sugar Sandwich Wagon C Baby Mankey Perfume Sunset Iron D Elbow Apple Carpet Saddle Bubble Jacket Pepper Cotton Movie Dollar Honey Mirror Saddle Anchor

DOB:		
Address:		
ID number:		
Examiner:		
Date:		

CONCENTRATION

DIGITS BACKWARDS

Please circle the Digit list chosen (A, B, C, D, E, F). Administer at the rate of one digit per second reading DOWN the selected column.

I am going to read a string of numbers and when I am done, you repeat them back to me in reverse order of how I read them to you. For example, if I say 7-1-9, you would say 9-1-7.

Concentra	tion Number Lis	ts (circle one)			
List A	List B	List C			
4-9-3	5-2-6	1-4-2	Y	N	0
6-2-9	4-1-5	6-5-8	Y	N	1
3-8-1-4	1-7-9-5	6-8-3-1	Y	N	0
3-2-7-9	4-9-6-8	3-4-8-1	Y	N	1
6-2-9-7-1	4-8-5-2-7	4-9-1-5-3	Y	N	0
1-5-2-8-6	6-1-8-4-3	6-8-2-5-1	Y	N	1
7-1-8-4-6-2	8-3-1-9-6-4	3-7-6-5-1-9	Y	N	0
5-3-9-1-4-8	7-2-4-8-5-6	9-2-6-5-1-4	Y	N	1
List D	Liet E	List F			
7-8-2	3-8-2	2-7-1	Y	N	0
9-2-6	5-1-8	4-7-9	Y	N	1
4-1-8-3	2-7-9-3	1-6-8-3	Y	N	0
9-7-2-3	2-1-6-9	3-9-2-4	Y	N	1
1-7-9-2-6	4-1-8-6-9	2-4-7-5-8	Y	N	0
4-1-7-5-2	9-4-1-7-5	8-3-9-6-4	Y	N	1
2-6-4-8-1-7	6-9-7-3-8-2	5-8-6-2-4-9	Y	N	0
8-4-1-9-3-5	4-2-7-9-3-8	3-1-7-8-2-6	Y	N	1
		Digits Score:			of 4

MONTHS IN REVERSE ORDER

Now tell me the months of the year in reverse order. Start with the last month and go be diward.

So you a say securious, restaurous, so arrest.	
Dec - Nov - Oct - Sept - Aug - Jul - Jun - May - Apr - Mar - Feb - Jan	0 1
Months Score	of 1
Concentration Total Score (Digits + Months)	of 5

Immediate Memory Score

Time that last trial was completed

STEP 4: NEUROLOGICAL SCREEN See the instruction sheet (page 7) for details of test administration and scoring of the tests. Can the patient read aloud (e.g. symptom check-list) and follow instructions without difficulty? Does the patient have a full range of pain-free PASSIVE cervical spine movement? Without moving their head or neck, can the patient look side-to-side and up-and-down without double vision? Can the patient perform the finger nose coordination test normally? Can the patient perform tandem gait normally? BALANCE EXAMINATION Modified Balance Error Scoring System (mBESS) testings ☐ Left ☐ Right Which foot was tested (i.e. which is the non-dominant foot) Testing surface (hard floor, field, etc.) Footwear (shoes, barefoot, braces, tape, etc.) _ Condition Errors of 10 Double leg stance Single leg stance (non-dominant foot) of 10 Tandem stance (non-dominant foot at the back) of 10 of 30 Total Errors

DOB:		
Address:		
ID number:		
Examiner:		

5		
STEP 5: D	ELAYED RECALI	L:
	he end of the Immediate	ed after 5 minutes have Recall section. Score 1
	at list of words I read a few time an remember in any order.	s earlier? Tell me as many words
	Time 5	itarted
Please record each w	ord correctly recalled. Total score	equals number of words recalled.
	_	
Total number of	words recalled accurately:	af5 or af10

6

STEP 6: DECISION

	Date & time of assessment:		
Domain			
Symptom number (of 22)			
Symptom severity score (of 132)			
Orientation (of 5)			
Immediate memory	of 15 of 30	of 15 of 30	of 15 of 30
Concentration (of 5)			
Neuro exam	Normal Abnormal	Normal Abnormal	Normal Abnormal
Balance errors (of 30)			
Delayed Recall	af 5 af 10	of 5 of 10	of 5 of 10

SCORING ON THE SCAT5 SHOULD NOT BE USED AS A STAND-ALONE METHOD TO DIAGNOSE CONCUSSION, MEASURE RECOVERY OR MAKE DECISIONS ABOUT AN ATHLETE'S READINESS TO RETURN TO COMPETITION AFTER CONCUSSION.

CLINICAL NOTES:	Name:
	DOB:
	Address:
	ID number:
	Date:
*	
CONCUSSION INJURY ADVICE	
(To be given to the person monitoring the concussed athlete)	Clinic phone number:
This patient has received an injury to the head. A careful medical examination has been carried out and no sign of any serious complications has been found. Recovery time is variable across	Patient's name:
individuals and the patient will need monitoring for a further pe-	Date / time of injury:
riod by a responsible adult. Your treating physician will provide guidance as to this timeframe.	Date / time of medical review:
If you notice any change in behaviour, vomiting, worsening head- ache, double vision or excessive drowsiness, please telephone your doctor or the nearest hospital emergency department immediately.	Healthcare Provider:
Other important points:	
Initial rest: Limit physical activity to routine daily activities (avoid exercise, training, sports) and limit activities such as school, work, and screen time to a level that does not worsen symptoms.	
1) Avoid alcohol	- Construction In Count Course 2017
Avoid prescription or non-prescription drugs without medical supervision. Specifically:	Concussion in Sport Group 2017
a) Avoid sleeping tablets	
 b) Do not use aspirin, anti-inflammatory medication or stronger pain medications such as narcotics 	
3) Do not drive until cleared by a healthcare professional.	
Return to play/sport requires clearance by a healthcare professional.	Contact details or stamp

INSTRUCTIONS

Words in Italics throughout the SCAT5 are the instructions given to the athlete by the clinician

Symptom Scale

The time frame for symptoms should be based on the type of test being administered. At baseline it is advantageous to assess how an athlete "typically" feels whereas during the acute/post-acute stage it is best to ask how the athlete feels at the time of testing.

The symptom scale should be completed by the athlete, not by the examiner. In situations where the symptom scale is being completed after exercise, it should be done in a resting state, generally by approximating his/her resting heart rate.

For total number of symptoms, maximum possible is 22 except immediately post injury, if sleep item is omitted, which then creates a maximum of 21.

For Symptom severity score, add all scores in table, maximum possible is 22 x 6 = 132, except immediately post injury if sleep item is omitted, which then creates a maximum of 21x6=126.

Immediate Memory

The Immediate Memory component can be completed using the traditional 5-word per trial list or, optionally, using 10-words per trial. The literature suggests that the immediate Memory has a notable ceiling effect when a 5-word list is used. In settings where this ceiling is prominent, the examiner may wish to make the task more difficult by incorporating two 5-word groups for a total of 10 words per trial. In this case, the maximum score per trial is 10 with a total trial maximum of 30.

Choose one of the word lists (either 5 or 10). Then perform 3 trials of immediate memory using this list.

elete all 3 trials regardless of score on previous trials

"I am going to test your memory, I will read you a list of words and when I am done. repeat back as many words as you can remember, in any order." The words must be read at a rate of one word per second.

Trials 2 & 3 MUST be completed regardless of score on trial 1 & 2.

Trials 2 & 3:

"I am going to repeat the same list again. Repeat back as many words as you can remember in any order, even if you said the word before.

ore 1 pt. for each correct response. Total score equals sum across all 3 trials. Do NOT inform the athlete that delayed recall will be tested.

Concentration

Digits backward

Choose one column of digits from lists A, B, C, D, E or F and administer those digits as follows:

Say: "I am going to read a string of numbers and when I am done, you repeat them back to me in reverse order of how I read them to you. For example, if I say 7:1-9, you would say 9-1-7."

Begin with first 3 digit string

If correct, circle "Y" for correct and go to next string length. If incorrect, circle "N" for the first string length and read trial 2 in the same string length. One point possible for each string length. Stop after incorrect on both trials (2 N's) in a string length. The digits should be read at the rate of one per second.

Months in reverse order

"Now tell me the months of the year in reverse order. Start with the last month and go backward. So you'll say December, November ... Go ahead*

1 pt. for entire sequence correct

Delayed Recall

The delayed recall should be performed after 5 minutes have elapsed since the end of the Immediate Recall section.

"Do you remember that list of words I read a few times earlier? Tell me as many words from the list as you can remember in any order.

Score 1 pt. for each correct response

Modified Balance Error Scoring System (mBESS)⁵ testing

This balance testing is based on a modified version of the Balance Error Scoring System (BESS)4. A timing device is required for this testing.

Each of 20-second trial/stance is scored by counting the number of errors. The examiner will begin counting errors only after the athlete has assumed the proper start position. The modified BESS is calculated by adding one error point for each error during the three 20-second tests. The maximum number of errors for any single condition is 10. If the athlete commits multiple errors simultaneously, only

one error is recorded but the athlete should quickly return to the testing position, and counting should resume once the athlete is set. Athletes that are unable to maintain the testing procedure for a minimum of five seconds at the start are assigned the highest possible score, ten, for that testing condition.

OPTION: For further assessment, the same 3 stances can be performed on a surface of medium density foam (e.g., approximately 50cm x 40cm x 6cm).

Balance testing - types of errors

- 1. Hands lifted off liac crest
- 3. Step, stumble, or fall 5. Lifting forefoot or heel
- 2. Opening eyes
- 4. Moving hip into > 30 degrees abduction
- 6. Remaining out of test position > 5 sec

"I am now going to test your balance. Please take your shoes off (If applicable), roll up your pant legs above ankle (if applicable), and remove any ankle taping (if applicable). This test will consist of three twenty second tests with different stances."

(a) Double leg stance:

"The first stance is standing with your feet together with your hands on your hips and with your eyes closed. You should try to maintain stability in that position for 20 seconds. I will be counting the number of times you move out of this position. I will start timing when you are set and have closed your eyes."

(b) Single leg stance:

"If you were to kick a ball, which foot would you use? [This will be the dominant foot] Now stand on your non-dominant foot. The dominant leg should be held in approximately 30 degrees of hip flexion and 45 degrees of knee flexion. Again, you should try to maintain stability for 20 seconds with your hands on your hips and your eyes closed. I will be counting the number of times you move out of this position. If you stumble out of this position, open your eyes and return to the start position and continue balancing. I will start timing when you are set and have closed your eyes."

"Now stand heel-to-toe with your non-dominant foot in back. Your weight should be evenly distributed across both feet. Again, you should try to maintain stability for 20 seconds with your hands on your hips and your eyes closed. I will be counting the number of times you move out of this position. If you stumble out of this position, open your eyes and return to the start position and continue balancing. I will start timing when you are set and have closed your eyes."

Tandem Gait

Participants are instructed to stand with their feet together behind a starting line (the test is best done with footwear removed). Then, they walk in a forward direction as quickly and as accurately as possible along a 38mm wide (sports tape), 3 metre line with an alternate foot heel to toe galt ensuring that they approximate their heel and toe on each step. Once they cross the end of the 3m line, they turn 180 degrees and return to the starting point using the same galt. Athletes fall the test if they step off the line, have a separation between their heel and toe, or if they touch or grab the examiner or an object.

Finger to Nose

"I am going to test your coordination now. Please sit comfortably on the chair with your eyes open and your arm (either right or left) outstretched (shoulder flexed to 90 degrees and elbow and fingers extended), pointing in front of you. When I give a start signal, I would like you to perform five successive finger to nose repetiti using your index finger to touch the tip of the nose, and then return to the starting position, as quickly and as accurately as possible."

References

- McCrory et al. Consensus Statement On Concussion in Sport The 5th nternational Conference On Concussion in Sport Held in Berlin, October 2016. British Journal of Sports Medicine 2017 (available at www.bjsm.bmj.com)
- Maddocks, DL; Dicker, GD; Saling, MM. The assessment of orientation following concussion in athletes. Clinical Journal of Sport Medicine 1995; 5: 32-33
- Jennett, B., Bond, M. Assessment of outcome after severe brain damage: a practical scale, Lancet 1975; I: 480-484
- McCrea M. Standardized mental status testing of acute concussion. Clinical Journal of Sport Medicine, 2001; 11: 176-181
- Guskiewicz KM. Assessment of postural stability following sport-related concussion, Current Sports Medicine Reports, 2003; 2: 24-30

C Concussion in Sport Group 2017

CONCUSSION INFORMATION

Any athlete suspected of having a concussion should be removed from play and seek medical evaluation.

Signs to watch for

Problems could arise over the first 24-48 hours. The athlete should not be left alone and must go to a hospital at once if they experience:

- Worsening headache
- Drowsiness or inability to be awakened
- Inability to recognize people or places
- · Repeated vomiting · Weakness or
- Unusual behaviour or confusion or irritable
- Seizures (arms and legs jerk uncontrollably)
- numbness in arms or legs
- Unsteadiness on their feet.
- · Slurred speech

Consult your physician or licensed healthcare professional after a suspected concussion. Remember, it is better to be safe.

Rest & Rehabilitation

After a concussion, the athlete should have physical rest and relative cognitive rest for a few days to allow their symptoms to improve. In most cases, after no more than a few days of rest, the athlete should gradually increase their daily activity level as long as their symptoms do not worsen. Once the athlete is able to complete their usual daily activities without concussion-related symptoms, the second step of the return to play/sport progression can be started. The athlete should not return to play/sport until their concussion-related symptoms have resolved and the athlete has successfully returned to full school/learning activities.

When returning to play/sport, the athlete should follow a stepwise, medically managed exercise progression, with increasing amounts of exercise. For example:

Graduated Return to Sport Strategy

Exercise step	Functional exercise at each step	Goal of each step
Symptom- limited activity	Daily activities that do not provoke symptoms.	Gradual reintroduc- tion of work/school activities.
2. Light aerobic exercise	Walking or stationary cycling at slow to medium pace. No resistance training.	Increase heart rate.
Sport-specific exercise	Running or skating drills. No head impact activities.	Add movement.
Non-contact training drills	Harder training drills, e.g., passing drills. May start progressive resistance training.	Exercise, coor- dination, and increased thinking.
5. Full contact practice	Following medical clear- ance, participate in normal training activities.	Restore confi- dence and assess functional skills by coaching staff.
6. Return to play/sport	Normal game play.	

In this example, it would be typical to have 24 hours (or longer) for each step of the progression. If any symptoms worsen while exercising, the athlete should go back to the previous step. Resistance training should be added only in the later stages (Stage 3 or 4 at the earliest).

Written clearance should be provided by a healthcare professional before return to play/sport as directed by local laws and regulations.

Graduated Return to School Strategy

Concussion may affect the ability to learn at school. The athlete may need to miss a few days of school after a concussion. When going back to school, some athletes may need to go back gradually and may need to have some changes made to their schedule so that concussion symptoms do not get worse. If a particular activity makes symptoms worse, then the athlete should stop that activity and rest until symptoms get better. To make sure that the athlete can get back to school without problems, it is important that the healthcare provider, parents, caregivers and teachers talk to each other so that everyone knows what the plan is for the athlete to go back to school.

Note: If mental activity does not cause any symptoms, the athlete may be able to skip step 2 and return to school part-time before doing school activities at home first.

Mental Activity	Activity at each step	Goal of each step
Daily activities that do not give the athlete symptoms	Typical activities that the athlete does during the day as long as they do not increase symptoms (e.g. reading, texting, screen time). Start with 5-15 minutes at a time and gradually build up.	Gradual return to typical activities.
2. School activities	Homework, reading or other cognitive activities outside of the classroom.	Increase tolerance to cognitive work.
3. Return to school part-time	Gradual introduction of school- work. May need to start with a partial school day or with increased breaks during the day.	Increase academic activities.
Return to school full-time	Gradually progress school activities until a full day can be tolerated.	Return to full academic activities and catch up on missed work.

If the athlete continues to have symptoms with mental activity, some other accomodations that can help with return to school may include:

- · Starting school later, only going for half days, or going only to certain classes
- More time to finish assignments/tests
- · Quiet room to finish assignments/tests
- Not going to noisy areas like the cafeteria, assembly halls, sporting events, music class, shop class, etc.
- · Taking lots of breaks during class, homework, tests
- · No more than one exam/day
- · Shorter assignments
- · Repetition/memory cues
- · Use of a student helper/tutor
- · Reassurance from teachers that the child will be supported while getting better

The athlete should not go back to sports until they are back to school/ learning, without symptoms getting significantly worse and no longer needing any changes to their schedule.

Appendix B Coaches/CMT Form



Coaches/CMT

Concussion Report

Time of Injury: Page Describe injury details:	arent Name/Phone:	Date of Injury:
Describe injury details: Any athlete who experience bump, blow, or jolt to the h		
 Any athlete who experience bump, blow, or jolt to the h 		
bump, blow, or jolt to the h		
	ead or body may have a con- lame. turn to play/practice until they	and symptoms listed below after a cussion and should be immediately have been evaluated by a health
Danger Signs: <u>If any are r</u>	resent, seek immediate i	medical attention, call 911
One pupil larger than the Repeated vomiting Slurred speech Convulsions or seizures	Cannot re Has unus	nsciousness ecognize people or places ual behavior nd cannot be awakened
symptoms Reported by Athlete	(Check all that apply)	
Headache or "pressure" in hea Nausea or vomiting Balance problems or dizziness Double or blurry vision Sensitivity to light/noise	Feeling Confus	not feel "right"
igns Observed by Coaching S	taff (Check all that apply)	
Appears dazed or stunned Forgets plays Moves clumsily Loses consciousness Is confused about plays	Can't r Answe Shows	ecall events prior to injury ecall events after injury rs questions slowly (days of the week etc.) behavior changes re of game, score, opponent
Completed by:	Signature	ə:
Contact parent/guardian of the in		

Regardless of whether the athlete is a key member of the team or the game is about to end, an athlete with a suspected concussion should immediately be removed from play. To help you know how to respond follow the CDC's "Heads Up" four-step action plan:

- 1. Remove the athlete from play.
- **2. Ensure** athlete is evaluated by appropriate health care professional.
- 3. Inform the athlete's parent or guardian.
- 4. Keep the athlete out of play.

Follow-up care instructions:

- o If any symptoms are getting worse, seek higher medical attention right away.
- Acetaminophen (Tylenol) is the only pain reliever that should be given for a concussion-related headache; avoid ibuprofen (Advil, Motrin) and aspirin.
- Gentle activity that doesn't worsen symptoms is encouraged after day 3 post concussion. The Center on Brain Injury Research and Training www.cbirt.org
- Athlete should also avoid TV, excessive reading, movies, computer use, tablet use, and texting since these activities will exacerbate the brain injury.
- Your athlete must be seen by a primary care physician or concussion specialist before returning to any physical activity.



To be evaluated immediately, check in through NOWcare at The Center Monday through Friday, 9 a.m. – 4 p.m.

If your athlete has a suspected concussion, he/she can be seen immediately through NOWcare at The Center located at 2200 NE Neff Road in Bend. Hours of operation are Monday through Friday 9 a.m. – 4 p.m. If you have any questions please contact Dr. Viviane Ugalde, Medical Director for concussion management at 541-322-2214.

For additional information on concussions, see below resources:

ImPACT Concussion Testing www.impacttest.com
The Center on Brain Injury Research and Training www.cbirt.org
CDC "Heads Up" www.cdc.gov/headsup
Brain 101 www.brain101.orcasinc.com

Appendix C Concussion Information Sheet



Parent Concussion Information Packet

Athlete Name:		Date:	
Date of Birth:	Date of Injury:	School:	

A concussion is a traumatic brain injury that alters the way a brain functions. Although concussions are usually caused by a blow to the head, they can also occur when the head and upper body are violently shaken causing the brain to be forced back and forth inside the skull. They can range from mild to severe and 90% of all concussions occur without a loss of consciousness. Signs and symptoms can present immediately or can take a few hours or days to fully appear. All concussions are potentially serious and may result in complications including prolonged brain damage and death if not recognized and managed properly.

Observed Signs & Symptoms

Pl	sical Thinking Emotional		Emotional	Sleep	
Headache	Sensitivity to light	Feeling mentally foggy	Irritable	Drowsiness	
Nausea	Sensitivity to noise	Problems concentrating	Sadness	Sleeping more than usual	
Fatigue	Numbness/tingling	Problems remembering	Feeling more emotional	Sleeping less than usual	
Visual problems	Vomiting	Feeling more slowed	Nervousness	Trouble falling asleep	
Dizziness	Balance problems				

RED FLAGS: Call your doo	tor or go to your emergen	cy department if you experience any of th	e following
Headache that worsens	Increasing confusion	Can't recognize people or places	Unusual behavior
Seizures	Repeated vomiting	Look very drowsy, can't be awakened	Increasing irritability
Slurred speech	Unequal pupils	Weakness/numbness in arms/legs	Loss of consciousness

If your athlete has a suspected concussion and is not exhibiting any of the above red flags, he/she should be seen by a health care professional within 24-72 hours. Your athlete can be seen immediately, without an appointment, through NOWCARE at The Center located at 2200 NE Neff Road in Bend. Hours of operation are M –F 9am – 4pm. Your athlete may also be seen by their primary care provider or through urgent care.

First 24 Hours after a Concussion: Common Questions

Q: Do I need to wake my child up every hour when s/he is sleeping?

A. No. Sleep is the best treatment for a concussion. It is OK to let him/her sleep without interruption the night of the injury after evaluation by a health care professional, or if you have spoken with your child's physician and s/he does not think your child needs further evaluation in the emergency department.

Q. Is it okay to give my child medicine for his/her headache?

A. Relieving headache pain is certainly appropriate, but it does not replace the need for cognitive and physical rest if symptoms are present. Be aware that symptom improvement with medication does not mean that the brain has recovered. After a concussion is diagnosed, talk to your physician about the use of medication — including type of medication and dose — for headache pain and other symptoms.

Q. My child wants to sleep all day long. Should I allow him/her to sleep as much as s/he wants?

A. A concussion affects how the brain works, so resting the brain as much as possible is necessary for recovery. Large amounts of sleep are normal. When your child is sleeping, his/her brain is recovering. It is a good idea to track the amount your child is sleeping and report it to your medical professional.

Q. Does my child need to give up sports if s/he has suffered a concussion?

A. Athletes should not return to sports while still having symptoms from a concussion because they are at risk for prolonging symptoms and further injury. It is very rare that any child is told to give up playing sports after a single injury. However, if the recovery is quite prolonged (greater than 6 months), you should consult with a concussion specialist to further discuss the possible risks of return to playing sports. An evaluation with a concussion specialist should be considered in any child who has had more than one sports-related concussion.

Q. My doctor told my child to have mental rest. What exactly does "rest" mean? Can my child watch television, play video games, text, etc.?

A. Mental rest means avoiding activities that require the brain to work hard to process information. This includes critical thinking and problem solving activities such as schoolwork, homework, and technology use. Restrictions from the following should be considered, because these activities increase brain function and can therefore worsen symptoms and delay recovery: Computer work/Internet use, video games, television, text messaging/cell phone use, bright lights, such as strobe lights at school dances, listening to loud music or music through headphones, loud noises, parties, concerts, pep rallies, driving, or work.

- See more at: cbirt.org/ocamp/parents

Returning to Daily Activities

- 1. Get lots of rest. Be sure to get enough sleep at night, try to keep the same bedtime.
- 2. Take daytime naps or rest breaks when you feel tired or fatigued.
- 3. Limit physical activity as well as activities that require a lot of thinking or concentration. These activities can make symptoms worse.
 - a. Physical activities include PE, sports practices, weight training, running, exercising, etc.
 - b. Thinking and concentration activities include homework, classroom work, job-related activity
- 4. Drink lots of fluids and eat carbohydrates and protein to maintain appropriate blood sugar levels
- 5. As symptoms decrease, you may begin to gradually return to your daily activities. If symptoms worsen or return, lessen your activities, then try again the next day to increase your activities gradually.
- 6. During recovery it is normal to feel frustrated and sad when you do not feel right and you can't be as active as usual.
- 7. Repeated evaluation of your symptoms is recommended to help guide recovery, see symptom tracking form.

Returning to School

- 1. After sustaining a concussion, if you awaken in the morning feeling poorly, headache, nauseated, dizzy, you should stay home from school and continue to rest as needed for the first three days.
- Extra help may be needed to perform school-related homework and classroom work, academic accommodations can be provided by physician to the school. These can be removed gradually as symptoms decrease.
- 3. If your symptoms linger and are not resolving or return, you should return to see the physician for further assessment.

Returning to Sports/Physical activity

- Once you are completely symptom free and attending school full time without academic accommodations, post-injury ImPACT testing will be completed. Those scores will be compared with your baseline, or if you do not have a baseline will be compared with age matched norms.
- 2. With approval from physician you may begin the graduated return to play steps under the supervision of athletic trainer, see Return to Play Post-Concussion Steps form.
- 3. Once all steps are completed without return of symptoms and written clearance is given by physician, you are cleared to participate in all physical activity.

Appendix D OSAA Concussion Return to Participation Form



Oregon School Activities Association 25200 SW Parkway Avenue, Suite 1 Wilsonville, OR 97070

503.682.6722 FAX 503.682.0960 http://www.osaa.org

CONCUSSION – RETURN TO PARTICIPATION MEDICAL RELEASE
Student Name: Date of Birth://_ School/Grade:
Date of Injury: / Sport/ Injury Details:
At this time, the student is: symptom-free at rest NOT symptom-free at rest NOT symptom-free at exertion NOT symptom-free at exertion Scoring within a normal range on ImPACT NOT scoring within a nor
Completed by (Printed name): Signature: Date:
Registered Athletic Trainer Coach Athletic Director Other:
Graduated, Step-wise Return-to-Participation Progression 1. No activity: Complete rest, both physical and cognitive. This may include staying home from school or limiting school hours and/or homework as activities requiring concentration and attention may worsen symptoms and delay recovery. 2. Light aerobic exercise: Walking or stationary bike at low intensity; no weight lifting or resistance training. Before progressing to the next stage the student must be healthy enough to return to school full time
 Sport-specific exercise: Sprinting, dribbling basketball or soccer; no helmet or equipment, no head impact activities. Non-contact training: More complex drills in full equipment. Weight training or resistance training may begin.
5. Full contact practice: Participate in normal training activities.
6. <u>Unrestricted Return-to-Participation/full competition</u> . (Earliest Date of Return-to-Participation:)
The student should spend a minimum of one day at each step. If symptoms re-occur, the student must stop the activity and contact their trainer or other health care professional. Depending upon the specific type and severity of the symptoms, the student may be told to rest for 24 hours and then resume activity one-step below where he or she was when the symptoms occurred. Graduated return applies to all activities including sports and PE classes.
This section to be completed by Physician/Health Care Professional:
Student may NOT return to any sport activity until medically cleared.
Student should remain home from school to rest and recover with a projected return date
Please allow classroom accommodations, such as extra time on tests, a quiet room to take tests, and a reduced workload when possible. Additional Recommendations:
Student may begin graduated return at stage circled above. If symptom free at rest and with graded exertion, can return to participation on date above.
Student is now cleared for full contact practice/participation: symptom free at rest and exertion and has completed a graduated Return-to-Participation protocol.
Physician/Health Care Professional Signature: Date:
Physician/Health Care Professional Name/Title:Phone:
Per OAR 581-022-0421 "Health Care Professional" means a Physician (MD), Physician's Assistant (PA), Doctor of Osteopathic (DO) licensed by the Oregon State Board of Medicine, nurse practitioner licensed by the Oregon State Board of Nursing, or Psychologist licensed by the Oregon Board of Psychologist Examiners.

Forms - Concussion-Return to Participation 07/11

2016-2017 Handbook



Oregon School Activities Association

25200 SW Parkway Avenue, Suite 1 Wilsonville, OR 97070

503.682.6722 FAX 503.682.0960 http://www.osaa.org

The Oregon School Activities Associations' (OSAA) Sports Medicine Advisory Committee has developed a physician release form for students to return to participation following a concussion. The committee reviewed extensively the literature available on concussions in sport. No definitive data exists that allow us to absolutely predict when a student with a concussion can safely return to participation. We have found significant differences that exist among physicians relating to when they will permit a student to return to participation after having a concussion.

Neither the OSAA nor the Sports Medicine Advisory Committee presumes to dictate to professionals how to practice medicine. Neither is the information on this form meant to establish a standard of care. The committee does feel, however, that the guidelines included on the form represent a summary consensus of the literature. The committee also feels that the components of the form are very relevant to addressing the concerns of coaches, parents, students, and physicians that lead to the research into this subject and to the development of this form.

GOALS FOR ESTABLISHING A WIDELY USED FORM:

- Protect students from further harm. Young students appear to be particularly vulnerable to the effects of concussion. They
 are more likely than older students to experience problems after concussion and often take longer to recover. Teenagers also
 appear to be more prone to a second injury to the brain that occurs while the brain is still healing from an initial concussion.
 This second impact can result in long-term impairment or even death. The importance of proper recognition and
 management of concussed young students cannot be over-emphasized.
- 2. Allow students to participate as soon as it is reasonably safe for them to do so.
- Establish guidelines to help minimize major differences in management among physicians who are signing "return to
 competition forms". Consistent use of these guidelines should minimize students from returning to participation too soon
 and protect them from inequalities as to who can or cannot participate.
- 4. Provide a basis to support physician decisions on when a student can or cannot participate. This should help the physician who may face incredible pressure from many fronts to return a student to competition ASAP. This can involve "Joe Blow who rides the bench" or the next state champion with a scholarship pending.

IMPORTANT COMPONENTS FOR AN EFFECTIVE FORM:

- Inclusion of the latest consensus statements so physicians will understand that students must be symptom free at rest and exertion and complete a graduated return to participation. Returning students at an arbitrary date is not an option.
- Inclusion of the date and nature of injury as well as earliest date to return to participation to minimize the need for a family to
 incur the expense of additional office visits to return for clearance after completing a graduated return to participation.
- Inclusion of consensus statements and return to participation progression before returning the student to participation as discussed above. This should enhance the likelihood that all students are managed safely and fairly.
- 4. Inclusion of all of the components discussed has the potential to remove liability from a school making a medical decision. If a return to participation is questioned, the school's role could appropriately be only to see if the student can provide a fully completed medical release form allowing the student to return to participation.

Note to Physicians/Health Care Professionals: Please familiarize yourself with the "Summary and Agreement Statements of International Conferences on Concussion in Sport", from Vienna in 2001, Prague in 2004, and Zurich in 2008. These documents summarize the most current research and treatment techniques in head injuries. The most noteworthy items to come from these conferences are the discontinuation of initial symptom based grading scales and the addition of standardized return to participation guidelines.

Note: ImPACT stands for Immediate Post-Concussion Assessment and Cognitive Test. It is sophisticated software developed to help sports-medicine clinicians evaluate recovery following concussion. ImPACT evaluates multiple aspects of neurocognitive functioning including memory, brain processing speed, reaction time, and post-concussive symptoms. For information on implementing a baseline-testing program, contact the Oregon Concussion Awareness & Management Program (OCAMP) at http://cbirt.org/ocamp.

Note: In 1990, the AMA recognized the certified athletic trainer as an allied health care professional. In 1998, a resolution passed urging all schools to provide the services of a certified athletic trainer for student-athletes (AMA Resolution 431, A-97). For more information on athletic trainers, contact Oregon Athletic Trainers' Society via their website: http://oatswebsite.org.

This form may be reproduced, if desired. In addition, the OSAA Sports Medicine Advisory Committee would welcome comments for inclusion in future versions, as this will continue to be a work in progress.

Forms - Concussion-Return to Participation 07/11

2016-2017 Handbook

Appendix E Return To Learn



Sports Concussion Program

Mild TBI/Concussion Temporary Accommodations Plan


Patient Name:	Date of Evaluation:
Current Symptoms: ☐ Hea ☐ Decreased attention ☐ Oth	daches Difficulty remembering Sensitivity to light Fatigue er:
	ecently suffered a concussion; symptoms and recovery timelines are individual to ecommendations and over time may need to be adjusted through the school
GENERAL RECOMMENDATIONS No return to school until specifi	i: ed. To be reviewed on:
 Shortened day:/_ No physical education classes w Physical education classes w May not be able to participa Consider reducing in-class w hours of nightly homework, 	other day, shortened day, shortened classes, breaks)hours/day
 Provide extended time to ta taking*) Allow breaks from electronic Provide written instructions 	emplete assignments and/or shortened assignments ke tests in a quiet environment (*do not mark if student is deferred from test cs if symptoms are present
 Allow rest breaks during the Allow "hall passing time" bet 	rse for treatment of headaches or other symptoms, if needed
athletic trainer	MOTIONAL ISSUES: progress and difficulties with parents, school nurse, counselor, physician and ort plan for the student (may include an adult with whom the student can talk,
Family has signed a release	e of information for bi-directional communication with
Signature:	Date:
Printed Name:	
Revised 10/23/15	

{000131470-00481657;3}

Appendix F Culver School District Concussion Policy Flowchart

Culver School District Concussion Protocol

All parents and athletes participating in sports have read and signed the Consent Form which includes Authorization for Use and Disclosure of Protected Health Information for Student Athletes and Concussion Acknowledgement (Family ID)



Appendix G Additional Resources

Additional resource available to coaches, teachers, and administrators

- Viviane Ugalde, MD: Medical Director of The Center Foundation Concussion Management Program, 541-382-3344
- St Charles Behavioral Health Pediatric Neuropsychologists 541-706-7730
- OCAMP (Oregon Concussion Awareness and Management Program) ocamp.org
- Brain 101 http://brain101.orcasinc.com/